

August 1994

**Wetland Status and Trends in
Charles County, Maryland
(1981 to 1988-89)**

U.S. Department of the Interior

Fish and Wildlife Service
Region 5



(Cowardin, *et. al.* 1979) following standard National Wetlands Inventory (NWI) mapping conventions (National Wetlands Inventory, 1990). These interpretations served as the basis for evaluating recent wetland trends.

The two sets of photographs were compared using a Bausch and Lomb SIS-95 zoom stereoscope. Changes were delineated on mylar overlays attached to the NAPP photographs. Cause of change was recorded for each polygon. The minimum mapping unit for wetlands was generally 0.5 acre, except for ponds, which were mapped when 0.1 acre or larger in size. Changes as small as 0.1 acre were detected. Wetland boundaries were improved and previously undetected wetlands were added to the original maps because the larger scale and more apparent signs of wetland hydrology of the NAPP photos improved our ability to detect and classify wetlands. Delineated changes and map refinements were then transferred to an NWI map using an Ottico Meccanica Italiana stereo facet plotter. Quality control of all photointerpretation was performed by a second photointerpreter. Tables were then prepared to present the study's findings.

RESULTS

Current Status

In 1988-89, Charles County contained about 27,010 acres of wetlands (roughly 9.3% of the County's land surface), excluding linear fringing wetlands along narrow streams. Table 1 summarizes the acreage of the different wetland types found in the County. Palustrine wetlands predominated, with 22,019 acres, representing 81.5% of the County's total wetland acreage. Nontidal deciduous forested wetlands accounted for 85.6% (18,859 acres) of all palustrine wetlands, and about 69.8% of the County's wetland total. Included within this total are significant portions of Zekiah Swamp, one of Maryland's largest freshwater wetlands. Tidal palustrine wetlands totaled 1,475 acres, representing 6.7% of the County's freshwater wetlands.

Estuarine wetlands comprised about 18.4% (4,969 acres) of the County's wetlands. Emergent wetlands (e.g., salt and brackish marshes) were the predominant type, accounting for almost 97% (4,804 acres) of the County's estuarine wetlands. These wetlands are located along tidal rivers and creeks emptying into Chesapeake Bay, and the Potomac, Patuxent, and Wicomico Rivers, among others. Slightly brackish marshes (oligohaline) are most common along Nanjemoy Creek, Port Tobacco Creek, Mattawoman Creek, and the Wicomico River (including Allens Fresh Marsh and Newport Marsh).

Recent Wetland Trends

Wetland trends results are presented in Tables 2 through 9. The following discussion highlights the more significant or interesting findings.

Vegetated Wetlands

Between 1981 and 1988-89, approximately 122 acres of vegetated wetlands were converted to upland (Table 2). Most of these losses affected palustrine forested wetlands. Housing construction was the most significant cause of vegetated wetland loss, with losses due to unknown factors also significant (Table 3). About 140 acres of vegetated wetland changed from one type to another. Upland conversion impacted the temporarily flooded palustrine wetland type more than others (Table 4). Approximately 292 acres of palustrine forested wetlands were converted to upland or changed to other wetland types (Table 5). Vegetated wetland gain from upland approached 48 acres (Table 6). Most gains in particular types of vegetated wetlands came from other vegetated wetland types (Table 6). Beaver activity affected 93 acres of vegetated wetlands, and created 39 acres of new wetlands by impounding upland areas (Table 7).

Nonvegetated Wetlands

About 105 acres of new ponds were created from upland, and close to 40 acres were constructed in vegetated wetlands (Table 8). More than 22 acres of ponds were converted to upland, while roughly 28 acres changed to vegetated wetlands. Approximately 27% of the new ponds built in uplands were constructed in urban areas, but the majority were attributed to other causes (Table 9).

CONCLUSION

The County had approximately 9.3% of its land mass covered by wetlands. Wetlands totaling 27,010 acres (in 1988-89) were identified in the County by the Service's National Wetlands Inventory. Palustrine wetland was the dominant type, representing 81.5% of the wetlands in the County.

Between 1981-82 and 1988-89, the County lost about 163 acres of vegetated wetlands, with roughly 122 acres converted to upland. Temporarily flooded wetland was the type most frequently converted to upland. Pond construction added about 135 acres of palustrine nonvegetated wetlands, but this gain was reduced to about 88 acres by pond losses to upland and vegetated wetlands.

The overall trend for the County's wetlands was losses of vegetated wetlands and gains in nonvegetated wetlands (mostly ponds). The significance of the increase in ponds to fish and wildlife species has not been assessed and remains a point for discussion. The losses of vegetated wetlands, however, represent known losses of valuable fish and wildlife habitats and areas providing other valued functions, including flood water storage, water quality enhancement, and local water supply.

While this report documents recent trends in the County's wetlands, it does not address changes in the quality of the remaining wetlands. As development increases, the quality of

wetlands can be expected to deteriorate due to agricultural runoff, increased sedimentation, groundwater withdrawals, increased water pollution, and other factors, unless adequate safeguards are taken to protect not only the existence of wetlands, but their quality.

ACKNOWLEDGMENTS

Funding for this project was provided by the Maryland Department of Natural Resources, Water Resources Administration through an existing cooperative agreement with the Service. David G. Burke was the project coordinator, and we appreciate his interest in monitoring wetland trends in Maryland.

Wetland maps and digital data were compiled by the U.S. Fish and Wildlife Service's National Wetlands Inventory Office at St. Petersburg, Florida. Special appreciation is extended to Becky Stanley and Linda Shaffer for their assistance. Photointerpretation was performed by the senior author and quality controlled by Glenn Smith. John Eaton compiled trend statistics, tables, and graphics for this report. Todd Nuerminger tabulated raw data.

REFERENCES

- Cowardin, L.M., V. Carter, F.C. Golet, and T. LaRoe. 1979. Classification of Wetlands and Deepwater Habitats of the United States. U.S. Fish and Wildlife Service, Washington, DC. FWS/OBS-79/31. 103 pp.
- Hoffman, M.S. (editor). 1992. The World Almanac and Book of Facts. Newspaper Enterprise Association, Inc., New York, NY. 960 pp.
- National Wetlands Inventory. 1990. Photointerpretation Conventions for the National Wetlands Inventory. U.S. Fish and Wildlife Service, St. Petersburg, FL. 45 pp. plus appendices.

Figure 1. Location of Study Area - Charles County, Maryland.




Table 1. Acreage of wetland types in Charles County, Maryland (1988-89).

<u>Wetland Type</u>	<u>Acres</u>	<u>% of Total</u>
PALUSTRINE WETLANDS		
Tidal Emergent		
Seasonally Flooded-Tidal	214.4	
Nontidal Emergent		
Semipermanently Flooded	102.7	
Seasonally Flooded/Saturated	154.3	
Seasonally Flooded	126.5	
Temporarily Flooded	109.1	
<i>(Subtotal Nontidal)</i>	<i>(492.6)</i>	1.8
Total Palustrine Emergent Wetlands	707.0	2.6
Tidal Forested		
Deciduous, Broad-leaved		
Semipermanently Flooded-Tidal	4.6	
Seasonally Flooded-Tidal	903.5	
Temporarily Flooded-Tidal	173.9	
Evergreen, Needle-leaved		
Seasonally Flooded-Tidal	4.4	
Temporarily Flooded-Tidal	1.7	
<i>(Subtotal Tidal)</i>	<i>(1,088.1)</i>	4.0
Nontidal Forested		
Evergreen, Needle-leaved		
Temporarily Flooded	59.3	
Seasonally Flooded	19.3	
Deciduous, Broad-leaved		
Seasonally Flooded/Saturated	1,741.9	
Seasonally Flooded	4,014.8	
Temporarily Flooded	12,774.5	
Semipermanently Flooded	324.8	
Dead	2.7	
<i>(Subtotal Nontidal)</i>	<i>(18,937.3)</i>	70.1
Total Palustrine Forested Wetlands	20,025.4	74.1

Table 1, continued

<u>Wetland Type</u>	<u>Acres</u>	<u>% of Total</u>
Tidal Scrub-Shrub	172.5	
Nontidal Scrub-Shrub		
Deciduous, Broad-leaved		
Seasonally Flooded/Saturated	168.0	
Seasonally Flooded	113.3	
Temporarily Flooded	143.0	
Semipermanently Flooded	30.5	
Permanently Flooded	1.0	
<i>(Subtotal Nontidal)</i>	<i>(455.8)</i>	1.7
Total Palustrine Scrub-Shrub Wetlands	628.3	2.3
Aquatic Bed	6.4	
Total Palustrine Vegetated Wetlands	21,367.1	79.1
Unconsolidated Bottom (Ponds)	626.9	
Unconsolidated Shore	25.1	
Total Palustrine Nonvegetated Wetlands	652.0	2.4
GRAND TOTAL PALUSTRINE WETLANDS	22,019.1	81.5
ESTUARINE WETLANDS		
Emergent		
Regularly Flooded	6.7	
Irregularly Flooded	1,178.5	
Regularly Flooded, Oligohaline	119.7	
Irregularly Flooded, Oligohaline	3,499.1	
Total Estuarine Emergent Wetlands	4,804.0	17.8

Table 1, continued

<u>Wetland Type</u>	<u>Acres</u>	<u>% of Total</u>
Scrub-Shrub		
Irregularly Flooded	23.1	
Irregularly Flooded, Oligohaline	82.7	
Total Estuarine Scrub-Shrub Wetlands	105.8	0.4
Forested, Irregularly Flooded	0.6	
Total Estuarine Forested Wetlands	0.6	
Total Estuarine Vegetated Wetlands	4,910.4	18.2
Total Estuarine Unconsolidated Shore	58.2	0.2
GRAND TOTAL ESTUARINE WETLANDS	4,968.6	18.4
RIVERINE WETLANDS		
Tidal Emergent	13.5	
Tidal Unconsolidated Shore	8.7	
GRAND TOTAL RIVERINE WETLANDS	22.2	0.1
TOTAL WETLANDS	27,009.9	100.0

Table 2. Changes of vegetated wetlands in Charles County, Maryland (1981 to 1988-89).

<u>Wetland Type</u>	<u>Converted to Upland (acres)</u>	<u>Changed to Other Vegetated Wetlands* (acres)</u>	<u>Changed to Nonvegetated Wetlands (acres)</u>	<u>Converted to Deepwater Habitat (acres)</u>
Palustrine Emergent	8.7	35.8	4.1	0.0
Palustrine Scrub-Shrub	5.8	16.4	0.3	0.0
Palustrine Forested	106.1	86.9	25.4	0.0
Estuarine Emergent	1.8	1.4	10.0	0.0
<u>Estuarine Scrub-Shrub</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>0.3</u>
Total	122.4	140.5	39.8	0.3

*Represents changes in wetland class (e.g., emergent to scrub-shrub) but not changes in water regime within a given wetland class.

Table 3. Causes of vegetated wetland loss to upland in Charles County, Maryland (1981 to 1988-89).

<u>Cause of Loss</u>	<u>Acres</u>
Housing Construction	44.9
Unknown Cause	30.9
Commercial Development	17.4
Road Construction	12.3
Agriculture	4.2
Industrial Development	1.9
<u>Ditching</u>	<u>1.1</u>
Total	122.6

Table 4. Conversion of hydrologically similar palustrine vegetated wetlands to upland in Charles County, Maryland (1981 to 1988-89).

<u>Palustrine Wetland Type</u>	<u>Acres</u>	<u>% Total Loss</u>
Temporarily Flooded	77.8	64.5
Seasonally Flooded	34.9	28.9
Seasonally Flooded/Saturated	1.1	0.9
Seasonally Flooded-Tidal	0.4	0.3
Semipermanently Flooded	1.7	1.4
<u>Permanently Flooded</u>	<u>4.9</u>	<u>4.0</u>
Total	120.8	100.0%

Table 5. Changes in palustrine forested wetlands in Charles County, Maryland (1981 to 1988-89).

<u>Forested Wetland Type</u>	<u>Converted to Upland (acres)</u>	<u>Changed to Other Wetland Types* (acres)</u>	<u>Total Loss (acres)</u>
Seasonally Flooded/Saturated	0.0	15.8	15.8
Seasonally Flooded	33.2	50.6	83.8
Temporarily Flooded	67.6	30.0	97.6
Semipermanently Flooded**	4.9	60.2	65.0
Seasonally Flooded-Tidal	0.4	23.4	23.8
<u>Temporarily Flooded-Tidal</u>	<u>0.0</u>	<u>5.7</u>	<u>5.7</u>
Total	106.1	185.7	291.7

*Includes both changes in wetland class (e.g., forested to emergent) and changes in water regime within a given wetland class.

**Represents dead forested wetlands.

Table 6. Gains in vegetated wetlands in Charles County, Maryland (1981 to 1988-89).

<u>Wetland Type</u>	<u>Gain from Nonvegetated Wetlands (acres)</u>	<u>Gain from Upland (acres)</u>	<u>Gain from Other Vegetated Wetlands (acres)*</u>
Palustrine Emergent	23.7	17.0	58.9
Palustrine Scrub-Shrub	2.1	3.2	65.4
Palustrine Forested	0.0	27.4**	0.0
Palustrine Aquatic Bed	2.0	0.0	12.2
<u>Estuarine Scrub-Shrub</u>	<u>0.0</u>	<u>0.0</u>	<u>4.0</u>
Total	27.8	47.6	140.5

*Represents changes in wetland class (e.g., emergent to scrub-shrub) but not changes in water regime within a given wetland class.

**Largely the result of beaver activity (all but 7.1 acres of this total, which were created as a result of man-made impoundments).

Table 7. Changes of wetlands in Charles County, Maryland due to beaver activity (1981 to 1988-89).

<u>Wetland Type</u>	<u>Change in Water Regime Only (acres)</u>	<u>Change in Vegetated Class (acres)</u>	<u>Gain from Upland (acres)</u>
Palustrine Emergent	12.1	2.4	3.3
Palustrine Scrub-Shrub	0.4	6.5	2.9
Palustrine Forested	63.3	8.7	20.3
Palustrine <u>Unconsolidated Bottom</u>	<u>0.0</u>	<u>0.0</u>	<u>12.5</u>
Total	75.8	17.6	39.0

Table 8. Gains and losses in nonvegetated wetlands in Charles County, Maryland (1981 to 1988-89).

Wetland Type	GAINS		LOSSES		
	Created from Upland (acres)	Created in Vegetated Wetlands (acres)	Converted to Upland (acres)	Changed to Vegetated Wetlands (acres)	Changed to Other Nonvegetated Wetlands (acres)
Palustrine					
Unconsolidated Bottom	104.8	39.7	18.2	17.5	0.3
Palustrine					
Unconsolidated Shore	0.0	0.0	0.0	10.3	0.0
Estuarine					
Unconsolidated Shore	<u>0.0</u>	<u>0.0</u>	<u>4.1</u>	<u>0.0</u>	<u>0.0</u>
Total	104.8	39.7	22.3	27.8	0.3

Table 9. Causes of recently constructed upland ponds in Charles County, Maryland (1981 to 1988-89).

<u>Causes</u>	<u>Pond Acreage</u>
Urban Ponds	28.2
Farm Ponds	25.4
Sand and Gravel Pit Ponds	19.2
Beaver Ponds	12.5
Other Ponds	9.6
Ponds in Undeveloped Areas	5.8
<u>Stormwater Detention Basins</u>	<u>4.1</u>
Total	104.8

